

Spectrum of Fatty Liver at Tertiary Care Centre of Northern India

Malhotra P*, Malhotra V, Sanwariya Y, GuptavU, Dixit, Girdhar S and Rathee S

Department of Medical Gastroenterology & Gynecology & Obstetrics, PGIMS, Rohtak and ADGHS, NVHCP, Haryana India

***Corresponding author:**

Parveen Malhotra,
Department of Medical Gastroenterology, Obstetrics & Gynecology and Microbiology PGIMS, Rohtak and ADGHS, NVHCP, Panchkula, 128/19, Civil Hospital Road, Rohtak, Haryana, 124001, India, E-mail: drparveenmalhotra@yahoo.com

Received: 25 Jun 2022

Accepted: 07 Jul 2022

Published: 13 Jul 2022

J Short Name: JJGH

Copyright:

©2022 Malhotra P, This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Keywords:

Fatty liver disease; Non-alcoholic fatty liver disease; Hepatitis B; Hepatitis C; Alcoholic liver disease

Citation:

Malhotra P. Spectrum of Fatty Liver at Tertiary Care Centre of Northern India. J Gastro Hepato. V9(1): 1-3

1. Abstract

1.1. Introduction: It is important to determine the epidemiological factors like etiology, age, sex, mode of clinical presentation of fatty liver and development of complications like chronic liver disease, so as to design optimal and cost effective preventive and treatment strategies for the same.

1.2. Aim: To determine spectrum of Fatty Liver at tertiary care centre of Northern India.

1.3. Material and Methods: This was a prospective study conducted at Medical Gastroenterology Department, PGIMS, Rohtak, over a period of five years i.e. 01.01.2017 to 31.12.2021, on 582 confirmed patients of Fatty Liver.

1.4. Results: The most common etiology seen was Non Alcoholic Fatty liver disease (82.47%), followed by Chronic HBV infection (11.37%), Chronic Hepatitis C (3.60%) and Alcoholic Liver Disease (2.57%),

1.5. Conclusion: The present study reveals that Nonalcoholic liver disease is most common cause of Fatty liver in Northern India.

2. Introduction

Fatty liver is the first insult to liver which can occur due to many reasons including Nonalcoholic Fatty liver disease (NAFLD), Alcoholic liver disease (ALD), Chronic Hepatitis B Virus (HBV) or Chronic Hepatitis C Virus (HCV) infection etc. The determination of fatty liver alarms the treating physician for further progression to hepatitis, cirrhosis and ultimately into Hepatocellular carcinoma (H.C.C.). NAFLD is a distinct hepatic disorder observed in patients without a history of significant alcohol consumption that histologically resem-

bles alcohol induced liver damage. It was first described in obese and diabetic women [1]. The spectrum of NAFLD passes from stage of simple Fatty liver to Non- alcoholic steatohepatitis (NASH), Cirrhosis and finally into small subset of patients into Hepatocellular carcinoma (H.C.C.) [2]. Like chronic hepatitis B, NAFLD in few cases, can directly progress to H.C.C without intervening stage of cirrhosis. Now universally NAFLD like Chronic HCV has become one of the most common etiological factor for causing end stage liver disease and requiring liver transplantation. NAFLD has been associated with insulin resistance, which includes obesity, diabetes, hypertriglyceridaemia and hypertension. In addition, NASH has also been associated with hyperlipoproteinemia, jejunal bypass, parenteral nutrition, drugs, α 1-antitrypsin deficiency, bacterial overgrowth, and environmental toxins. Exposure to chemicals is suspected to be another risk factor. Iron overload, H63D mutation of the HFE gene, and immune system anomalies are frequent in patients with non- alcoholic steatohepatitis. Recent reports have also described hepatocellular carcinoma arising in patients with NASH associated cirrhosis [3]. Chronic HBV and HCV infection usually remain asymptomatic for a long period and thus impact on liver which begins with fat infiltration is not detected in many. A certain subset of patient in chronic HBV and HCV infection present in late stage i.e. cirrhosis. Thus in these patients fatty liver stage has already progressed to cirrhotic stage. In majority of cases of ALD, the consultation is sought at later advanced stage when already cirrhosis has set in. Thus fatty liver stage is not frequently seen. The ALD patients are usually reluctant for timely consultation, as they know that once they are diagnosed to be having side effects of alcohol on their body, then it will lead to pressure on leaving alcohol from family and friends.

3. Aim

To determine spectrum of Fatty Liver at tertiary care centre of Northern India.

4. Material and Methods

This was a prospective study done at Medical Gastroenterology Department, PGIMS, Rohtak conducted over a period of five years i.e. 01.01.2017 to 31.12.2022, on 582 confirmed cases of Fatty liver. The patients who visited the Medical Gastroenterology Department in last five years and were confirmed to be having only Fatty Liver on ultrasonogram and no cirrhotic changes, after proper consent were enrolled in the study. Their detailed records were collected regarding aetiological and epidemiological factors, clinical spectrum and were followed regularly. The detailed clinical examination and laboratory investigations were done including complete blood counts, liver function tests, kidney function tests, thyroid function test, serum IgA TIG antibody, serum electrolytes, coagulation parameters (PT, INR), blood sugar, autoimmune profile, Serum ceruloplasmin level, 24 hour urinary copper excretion level, Hbs Ag, anti HIV antibody, anti HCV antibody, HBV DNA Quantitative, HCV RNA Quantitative, ultrasonogram abdomen, chest x ray PA view and Fibroscan.

5. Observations

Out of the total study group of 582 confirmed patients of Fatty liver, the most common etiology determined was Non Alcoholic liver disease (480 patients, 82.47%), followed by Chronic Hepatitis B (66 patients, 11.37%), Chronic Hepatitis C (21 patients, 3.60%) and Alcoholic liver disease (15 patients, 2.57%).

On individual analysis of Non Alcoholic Fatty liver disease, out of total 480 patients, 228 (47.50%) were male and 252 patients (52.50%) were female. On analysis of rural-urban distribution, 282 patients (58.75%) belonged to rural areas and 198 patients (41.25%) belonged to urban areas. On analyzing age distribution curve, none patient was seen in 0-10 yrs of age after which gradual upward trend was noted i.e. 11-20 years (15 patients, 3.12%), 21-30 yrs (72 patients, 15%) and peak was seen between 31-40 yrs (126 patients, 22.10%), followed by gradual decline i.e. 41-50 yrs (102 patients, 21.25%), 51-60 yrs (90 patients, 18.75%), 61-70 yrs (66 patients, 14.37%), 71-80 yrs (1 patient, 0.6%) and 81-90 yrs (1 patient, 0.6%). Out of total 480 patients, a significant proportion i.e. 174 patients, 36.25% were in stage of Nonalcoholic steatohepatitis as evidenced by raised transaminases. The Dyslipidemia was seen in 120 patients (25%), 36 patients (12.50%) were diabetic and 24 patients (5%) were hypothyroid.

On individual analysis of 66 patients (11.34%) of Chronic Hepatitis B related Fatty liver, 51 (77.27%) were male and 15 patients (22.73%) were female, 54 patients (81.81%) belonged to rural areas and 12 patients (18.19%) belonged to urban areas. On analyzing age distribution curve, none patient was seen in 0-10 yrs of age after which gradual upward trend was noted i.e. 11-20 years (3 patients, 4.54%), 21-30 yrs (12 patients, 18.18%) and peak was seen between 31-40 yrs (21 patients, 31.81%), followed by gradual decline i.e. 41-50 yrs (12

patients, 18.18%), 51-60 yrs (6 patients, 9.09%), 61-70 yrs (12 patients, 18.18%), 71-80 yrs (1 patient).

The analysis of 21 patients (3.60%) of Chronic hepatitis C related Fatty liver group, 18 (85.71%) were male and 3 patients (14.29%) were female, 18 (85.71%) belonged to rural areas and 3 patients (14.29%) belonged to urban areas. On analyzing age distribution curve, none patient was seen in 0-30 yrs of age after which gradual upward trend was noted i.e. 31-40 yrs (6 patients, 28.57%), 41-50 yrs (12 patients, 57.14%) and 51-60 yrs (3 patients, 14.28%).

On analyzing 15 patients, 2.57% of Alcoholic Fatty liver disease group, all 15 were males (100%), 12 patients (80%) belonged to rural areas and 3 patients (20%) belonged to urban areas. On analyzing age distribution curve, none patient was seen in 0-30 yrs of age after which gradual upward trend was noted i.e. 31-40 yrs (6 patients, 40%), 41-50 yrs (3 patients, 20%) and 51-60 yrs (6 patients, 40%) (Table 1-5).

Table 1: Showing Distribution of Fatty Liver Patients on Basis of Etiology

Category	Number of Patients (582)
NAFLD	480 (82.47%)
HBV	66 (11.34%)
ALD	21 (3.60%)
HCV	15 (2.57%)

Table 2: Showing Sex Distribution of Fatty Liver Patients.

Sex	NAFLD (480)	HBV (66)	HCV (21)	ALD (15)
Male	228 (47.50%)	51 (77.27%)	18 (85.71%)	15 (100%)
Female	252 (52.50%)	15 (22.73%)	3 (14.29%)	0 (0%)

Table 3: Showing Rural and Urban Distribution of Fatty Liver Patients

Residence	NAFLD (480)	HBV (66)	HCV (21)	ALD (15)
Rural	282 (58.75%)	54 (81.81%)	18 (85.71%)	12 (80%)
Urban	198 (41.25%)	12 (18.19%)	3 (14.29%)	3 (20%)

Table 4: Showing Age Distribution of Fatty Liver Patients

Age Group	NAFLD (480)	HBV (66)	HCV (21)	ALD (15)
0-10	0	0	0	0
20-Nov	15	3	0	0
21-30	72	12	0	0
31-40	126	21	6	6
41-50	102	12	12	3
51-60	90	6	3	6
61-70	69	12	0	0
71-80	3	0	0	0
81-90	3	0	0	0

Table 5: Showing Distribution of Parameters NAFLD Group

NAFLD	NASH	Dyslipidemia	Diabetic	Hypothyroid
480 Patients	174 patients	120 patients	36 patients	24 patients
Percentage	36.25%	25%	12.50%	5%

6. Discussion

There are many studies worldwide on NAFLD but there is paucity of data for other etiologies and their contribution in total pool of Fatty liver disease. This was the reason for conducting this study which was hospital based study with aim of determining the etiology of Fatty liver disease in our region. Male predominance was seen in all the various groups of Fatty liver diseases except in NAFLD group. The majority of patients in all the groups were seen in middle age group (30-60 yrs). In comparison to other etiologies like HBV, HCV and ALD, NAFLD patients were seen at both extreme of ages i.e. from 10 yrs to 85 yrs. We know that Paediatric NAFLD has assumed significant percentage in overall NAFLD group worldwide. The NAFLD in elderly can be explained on basis of occurrence at later ages and slower progression in comparison to other etiologies. That means HBV, HCV and ALD patients either recover due to stopping alcohol or by treatment or succumb to their illness before reaching to much elder age group. If comparison is made with etiological reasons for chronic liver disease, as shown by Malhotra et al [4], in which most common cause found was Alcoholic liver disease (48.9%), followed by Nonalcoholic liver disease (26.4%), Chronic Hepatitis B (12.3%), Chronic Hepatitis C (9%), Cryptogenic (2.7%) and Autoimmune liver disease related (0.7%) which is in contrast to Fatty liver disease spectrum in which NAFLD predominates. The reason behind this can be that majority of ALD patients present late as they are averse to take treatment for the same. Regarding HBV infection explanation can be that majority of new patients diagnosed are incidentally detected, are asymptomatic and have low viral load i.e. are chronic inactive carriers and only few are in active stage which can manifest as normal liver on ultrasonogram, Fatty liver or cirrhotic changes. In HCV also, low percentage of Fatty liver seen can be due to incidental detection in majority of patients who are asymptomatic. In our study, rural predominance was seen in all the groups of Fatty liver disease. Out of total pool of 480 patients in NAFLD group, steatohepatitis as evidenced by raised transaminases was seen in 174 patients (36.25%). The Dyslipidemia was seen in 120 patients (25%) and in this most common was increased cholesterol level followed by increased Serum triglycerides and Low density lipoprotein (LDL) levels. In NAFLD group, 36 patients (12.50%) were diabetic and 24 patients (5%) were hypothyroid. NAFLD has become second most common reason for Chronic Liver Disease (CLD) in India because of increasing trend of sedentary life style and contribution from increased number of patients of Diabetes Mellitus. In India, under National Viral Hepatitis Control Program (NVHCP), free treatment is being given for hepatitis B & C all over India for curbing the menace of these deadly diseases. As per results of our research, there is urgent need of taking steps for determining and controlling

the factors leading to alcoholic liver disease and Nonalcoholic fatty liver disease because of their significant contribution in total pool of CLD in India. In both of these etiological factors, public awareness on large scale is required because it is well said that prevention is better than cure. Moreover, Liver transplant which is the only definitive treatment for End Stage liver disease, is beyond reach of majority of Indians due to its limited availability, less number of organ donation and financial constraints. Hence, in developing country like India, aim should be not to make patients reach End stage liver disease and on transplant waitlist.

7. Conclusion

Fatty liver disease is common entity in Northern India with male preponderance except in NAFLD and affecting mostly people of middle age group. NAFLD is major cause of Fatty liver disease, followed by Chronic Hepatitis B & C and ALD. The casual approach of nothing to worry for Fatty liver should be changed to aggressively dealing with the etiology causing Fatty liver, so as to prevent its progression to hepatitis, cirrhosis and H.C.C. Hence government should prioritize on making national health policy on NAFLD as done for hepatitis B & C under National viral hepatitis control program (NVHCP).

References

1. Ludwig J, Viggiano TR, McGill DB, Ott BJ. Nonalcoholic steatohepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin Proc.* 1980; 55: 434-8.
2. Das SK, Vasudevan DM. Drugs and non alcoholic steatohepatitis. *Ind J Pharmacol.* 2006; 38(4): 238-242
3. Das SK, Mukherjee S, Vasudevan DM. Non-alcoholic fatty liver diseases: an under recognized cause with emerging importance. *Curr Sci.* 2006; 90(5): 659-665.
4. Malhotra P, Malhotra V, Sanwariya Y, Chugh A, Pahuja I, Akshay. Spectrum of Chronic Liver Diseases in India. *Adv Res Gastroenterol Hepatol* 2021; 17(3): ARGH.MS.ID.555964.